

# Effective interactions for reaction calculations? A first look at the effect of density dependence

**Jutta Escher, F.S. Dietrich, G.P.A. Nobre, I.J. Thompson**  
Nuclear Theory & Modeling, LLNL

**M. Dupuis**  
LANL

**A.K. Kerman**  
ORNL/MIT

**Lawrence  
Livermore National  
Laboratory**



**SciDAC/UNEDF Annual  
Collaboration Meeting  
Pack Forest, WA  
June 23-26, 2008**

# UNEDF Reaction Theory

**Goal:** Replace phenomenological input for reaction calculations by theoretically predicted quantities in order to enable reaction calculations for areas with little/no available systematics.

## Strategy:

- Employ state-of-the-art nuclear theory input (e.g. nuclear structure input from DFT)
- Employ large-scale computational resources to minimize/eliminate those inputs that lump together physics that cannot be described properly with traditional tools (e.g. “channels”)

## Practical approach:

- Implement a large-scale coupled-channels framework that is able to exploit the available (predicted) nuclear structure input.
- Develop/determine and test the ingredients that cannot be supplied by DFT.

### Coupled-channels equations:

$$\left\{ E - E_c^{\text{ex}} + \frac{\hbar^2}{2\mu_c} \frac{d^2}{dr^2} - \frac{l_c(l_c+1)\hbar^2}{2\mu_c r^2} - V_{cc}^{\text{eff}}(r) \right\} u_c(r) = \sum_{c' \neq c} V_{cc'}^{\text{eff}}(r) u_{c'}(r),$$

The coupling potentials  $V_{cc'}^J(r)$  contain information on the structure of the target nucleus and on the effective NN interaction between the projectile (nucleon) and target nucleons, via terms of the type:

$$\frac{2}{\pi} \int_0^\infty dq q^2 j_L(qr) v^{ST}(q) \rho_{LSJ}^{Tq, \tilde{n}}(q)$$

Transition densities for target states, obtained from a nuclear-structure calculation based on DFT results.

Effective NN interaction between a particle in the continuum and a nucleon in the target - to be determined!



# Which effective interaction?

Coupling potentials require folding of effective interaction with transition densities (in r or q space):

$$\frac{2}{\pi} \int_0^{\infty} dq q^2 j_L(qr) v^{ST}(q) \rho_{LSJ}^{Tq, \mathbf{f}}(q)$$

It is not obvious which effective interaction to use for calculating the coupling potentials.....

Some of these issues have been considered in the context of the OMP....  
....for the diagonal term.

The OMP is the potential that reproduces the observed elastic scattering cross section when used in a one-channel model.

Questions:

- Which effective interaction?
- Should we use the same that was employed for the structure calculation?
- How do we account for the fact that one of the interacting particles is in the continuum?
- How do we connect with a Skyrme functional?
- Does it matter which (fitted) interaction we use?
- If so, can we identify constraints for selecting a good effective interaction?




# Which effective interaction? Insights from OMPs

Prior considerations in the context of the “standard” OMP:

- Single- and double-folding approach commonly used to determine the real portion of an optical potential for standard reaction calculations.
- **Density dependence** is important for describing inelastic  $\alpha$ -nucleus scattering.
- **Energy dependence**: Phenomenological OMPs have been extrapolated to the  $E < 0$  region to calculate bound single-particle states.

**These issues have NOT been studied in the context of coupled-channels calculations that explicitly include a very large number of channels.**

OMP description of elastic scattering with OMPs obtained from folding with effective interactions with different density dependencies



From: Khoa et al, J. Phys. G 34 (2007) R111



# Density dependence of the effective interaction

## Constructing diagonal coupling potentials from effective interactions with/without density dependence:

- Effective nucleon-nucleon interaction:

$$V(E, \rho, r) = F(\rho) v(r)$$


Where  $v$  is a M3Y interaction

$$v(r) = \sum_{i=1..3} G_i \exp(-\mu_i r) / (\mu_i r) + J_{00}$$


$$F(\rho) = C(1 + \alpha \exp(-\beta \rho))$$

$$J_{00} = -G_\delta (1 - 0.0005 E/A)$$

- Density from a Skyrme (SkM\*) HF calculation






Coupling potentials compared



# Constraints from single-particle energies?

Neutron single-particle states in  $^{90}\text{Zr}$



The folded (diagonal) coupling potential can be used to predict single-particle energies for bound orbitals. These can be compared to data and to DFT results.




# Effects on coupled-channels results

Coupled-channels calculation of reaction cross section for  $n+^{90}\text{Zr}$ , employing different effective interactions in the diagonal coupling potential.

-> Results for density-dependent interaction improve agreement with data.

See also G. Nobre's presentation.




# Effect on DWBA inelastic scattering results

DWBA calculation of inelastic n scattering cross section, employing different effective interactions in the (off-diagonal) transition potentials

-> Illustrates the impact of not having a definite prescription for selecting an effective interaction.

See also M. Dupuis' presentation.



# Plan for Year 2 and Year 3

---

## Determine effective interaction appropriate for reaction calculations

- Implement folding procedure for various candidate effective interactions (M3Y, DDM3Y, CDM3Yx, Gogny, Gogny at  $\rho=0$ , others) Done for select cases.  
Others to be considered.
- Calculate the diagonal ( $c=c'$ ) potential and investigate effects of varying the effective forces with special emphasis on density and energy dependence Started.
- Devise tests to constrain the coupling potentials and thus the effective interactions:  
– Single-particle energies for the diagonal potential  
– Energy-dependence must satisfy the dispersion relation  
– Positions of doorway resonances, etc. Started.  
Needs to be expanded.
- Determine sensitivity of cross sections to choice of effective interaction Started.
- Draw conclusions about the possibility of moving from an effective interaction for bound states to an interactions valid for positive energies To be done.
- Recommend interaction(s) for reaction calculations (all couplings). To be done.



# APPENDIX

---



# Effective nucleon-nucleon interaction

## Effective nucleon-nucleon interaction - M3Y with $\rho$ -dependence

$$V(E, \rho, r) = g(E) F(\rho) v(r)$$

where

$$v(r) = \sum_{i=1..3} G_i \exp(-\mu_i r) / (\mu_i r) + J_{00}$$

$$F(\rho) = C(1 + \alpha \exp(-\beta\rho) - \gamma\rho)$$

$$g(E) = 1 - k_p E$$



$$J_{00} = -G_\delta(1 - 0.0005E/A)$$

F( $\rho$ )( $\gamma=0$ )	Reid	Paris
C	0.2845	.2963
$\alpha$	3.6391	3.7231
$\beta$ [fm <sup>3</sup> ]	2.9605	3.7384

Coefficient	Reid	Paris
$G_1$ [MeV]	7999	11062
$G_2$ [MeV]	-2134	-2537.5
$G_3$ [MeV]	0	0
$\mu_1$ [fm <sup>-1</sup> ]	4	4
$\mu_2$ [fm <sup>-1</sup> ]	2.5	2.5
$\mu_3$ [fm <sup>-1</sup> ]	0.7072	0.7072
$G_\delta$ [MeVfm <sup>3</sup> ]	-276	-592
k [MeV <sup>-1</sup> ]	0.002	0.003



# Densities and mean field from Skyrme



# In context...

